



INDICATOR: GROUND-LEVEL OZONE

STRATEGIC DIRECTION: Reduce Threats

TARGET: N/A

THEME: Pressures on Ontario's Biodiversity

Background Information:

Ground-level ozone is a secondary pollutant, formed by chemical reactions between nitrogen oxides and volatile organic compounds in the presence of heat and sunlight. Nitrogen oxides are emitted by natural sources and by human sources such as cars, trucks, and industrial plants. Volatile organic compounds may be emitted by natural sources, such as plants and trees, or by human activities. In addition to having impacts on human health, there is substantial evidence that ground-level ozone can have significant impacts on biodiversity. The most important and well documented impacts are reported for terrestrial vegetation with reductions in crop production, tree growth, and changes in species composition observed (Hayes et al. 2007; Wittig et al. 2009), all of which have downstream consequences for ecosystem services (Royal Society 2008). Evidence suggests that these impacts are likely to occur where ozone concentrations are above a threshold of 40 ppb, a limit which is currently exceeded in most countries around the world, including Canada (Royal Society 2008; Environment Canada 2014).

This indicator is used to assess trends in the seasonal means of ground-level ozone at sites across Ontario, as well as the annual peak (4th-highest) daily maximum 8-hour concentration.

Data Analysis:

Data to assess trends in the seasonal means of ground-level ozone concentrations in Ontario from 1980 to 2012 (Fig. 1) were obtained from the Air Quality in Ontario Reports 2007 and 2012 (OMOE2008; OMOECC 2014). Seasonal means were based on data from several ozone monitoring stations operated across Ontario. A 75% data completeness criterion was used to derive trends. Ozone long-term trends were derived from 19 sites for 1980 to 2007 (OMOE 2008) and from 37 sites from 2003 to 2012 (OMOECC 2014). The stations were largely based in metropolitan areas in Ontario.

Data used to assess annual peaks (4th highest) for daily maximum 8-hour ground-level ozone concentration from 1998 to 2012 (Fig. 2) were obtained from the Canadian Environmental Sustainability Indicator Report 2014 (Environment Canada 2014) which includes data from National Air Pollution Surveillance Program (NAPS) and the Canadian Air and Precipitation Monitoring Network (CAPMoN). The annual peaks for ground-level ozone are calculated using an approach that is aligned with the Canadian Ambient Air Quality Standards (CAAQS). Annual peaks are based on data collected from 34 monitoring stations in southern Ontario. Peaks are based on the 4th-highest of the daily maximum 8-hour average concentrations measured over a given year. Annual peaks for Ontario were obtained for this indicator by averaging all the 4th-highest values from stations in this region.

- [Link to Environment Canada Data](#)



Results:

Trend: Mixed **Data Confidence:** High **Geographic Extent:** Provincial




Figure 1. Seasonal means of ground-level ozone at sites across Ontario 1980-2012 (Source: OMOECC 2008 and 2014).




Figure 2. Annual peak (4th-highest) daily maximum 8-hour ground-level ozone concentration at sites across Ontario 1998-2012 ($n = 34$) (Source: Environment Canada 2014) (Note: The red horizontal dashed line represents the value of the Canadian Ambient Air Quality Standards (CAAQS) for 2015 and is shown for indicative purposes only, and not for evaluation of the achievement status of the standard. The green horizontal dashed line represents the environmental threshold above which impacts are likely on vegetation (Royal Society 2008)).



Status:

- Between 1980 and 2012 seasonal means of ground-level ozone in Ontario increased in both the summer and winter seasons. However, during the past 10 years the trend in summer means has remained constant, while the winter means increased by 16% over the same 10-year period. The increase in ozone winter means is mainly attributed to rising global background concentrations.
- A decreasing trend was detected in the annual peak concentrations of ground-level ozone from 1998 to 2012, representing a decrease of 17% (or an average of 1.2% per year) over that period. A reduction in emissions of ground-level ozone precursor gases (nitrogen oxides and volatile organic carbons) from Canada and the United States is likely an important factor in this downward trend. However, the annual peak ground-level ozone concentrations were above the 2015 CAAQS and environmental threshold for all years.

Links:

Related Targets: 8. By 2015, the release of pollutants harmful to biodiversity is reduced.

Related Themes: N/A

Web Links:

Canadian Environmental Sustainability Indicators <http://www.ec.gc.ca/indicateurs-indicators/default.asp?lang=En>

Ontario Ministry of Environment and Climate Change Air Quality Reports <http://www.airqualityontario.com/press/publications.php>

National Air Pollution Surveillance Program (NAPS) <http://www.ec.gc.ca/rnspace-naps/>

Canadian Air and Precipitation Monitoring Network (CAPMoN) <https://www.ec.gc.ca/rs-mn/default.asp?lang=En&n=752CE271-1>

Canadian Ambient Air Quality Standards (CAAQS) <http://www.ec.gc.ca/default.asp?lang=En&n=56D4043B-1&news=A4B2C28A-2DFB-4BF4-8777-ADF29B4360BD>

References:

Environment Canada. 2014. Canadian Environmental Sustainability Indicators. [Available at: <https://www.ec.gc.ca/indicateurs-indicators/default.asp?lang=en&n=7DCC2250-1>.]

Hayes F., M.L.M. Jones, G. Mills, and M. Ashmore. 2007. Meta-analysis of the relative sensitivity of semi-natural vegetation species to ozone. *Environmental Pollution* 146:754-762.

Lovett, G.M., T.H. Tear, D.C. Evers, S.E.G. Findlay, B.J. Cosby, J.K. Dunscomb, C.T. Driscoll, and K.C. Weathers. 2009. Effects of air pollution on ecosystems and biological diversity in the eastern United States. *BioScience* 1162:99-135.

Ontario Ministry of the Environment and Climate Change (OMOECC). 2014. Air quality in Ontario report 2012. Queen's Printer for Ontario, Toronto, ON.

Ontario Ministry of the Environment (OMOE). 2008. Air quality in Ontario Report 2007. Queen's Printer for Ontario, Toronto, ON.

Royal Society. 2008. Ground-level ozone in the 21st century: future trends, impacts and policy implications. Science Report, Royal Society, London.



Wittig V.E, E.A. Ainsworth, S.L. Naidu, D.F. Karnosky, and S.P. Long. 2009. Quantifying the impact of current and future tropospheric ozone on tree biomass, growth, physiology and biochemistry: a quantitative meta-analysis. *Global Change Biology* 15:396-424.

Citation

Ontario Biodiversity Council. 2015. State of Ontario's Biodiversity [web application]. Ontario Biodiversity Council, Peterborough, Ontario. [Available at: <http://ontariobiodiversitycouncil.ca/sobr> (Date Accessed: May 19, 2015)].